# GENERAL INDEX TO VOLUME XVIII

New scientific names of plants and the final members of new combinations are printed in **bold face** type; synonyms, and page numbers having reference to figures and plates, in *italics;* and previously published names and all other matter, in ordinary type

---


## A

abietinus (Polystictus), 470
Abietis (Trametes), 474
Acidity of Missouri soils and its relation to plant distribution, A study of, 41
Adansonia, 128
adustus (Polyporus), 471
Agaricaceae, 470
*Agaricus boletiformis*, 391; *hirsutus*, 391; *saepiarius*, 391; *trabeus*, 392
Aleurodiscus disciformis, 485; **diffissus**, 485
Allium stellatum, Chromosome complements of, 465, *468;* Moly, 465; ursinum, 465
**Alpova**, a new genus of Rhizopogonaceae, with further notes on Leucogaster and Arcangeliella, 457
**Alpova cinnamomeus**, 461, *464*
Althaea, 118, 125
ambiens (Helicoma), 10
*ammonis* (*Gyroceras*), 1, 7
**ammonis** (Sarcopodium) (Tricholeconium), 7
amorphus (Polyporus), 471
**anastomosans** (Helicoma), 13, 14, *16*
Anderson, Edgar. The chromosome complements of Allium stellatum and Nothoscordum bivalve, 465; and Dorothea De Winton. The genetic analysis of an unusual relationship between self-sterility and self-fertility in Nicotiana, 97
*Anechites adglutinata*, 552
*Anisolobus cognatus*, 546; *Perrottetii*, 546
*Antrodia mollis*, 367
Apocynaceae of tropical America, New or otherwise noteworthy, 541
applicatus (Pleurotus), 470
Arcangeliella **australiensis**, 463; **Ravenelii**, 463; **rosea**, 462; **Stephensii**, 463
Arenaria cephaloidea, 284
arida (Coniophora), 485
Asclepiadaceae, New South American, 557
Asia, eastern, Hymenomycetous fungi of, 469
*Aurantiporellus alboluteus*, 333
auricula-Judae (Auricularia), 487
Auricularia auricula-Judae, 487; auriformis, 487
Auriculariaceae, 487
auriformis (Auricularia), 487
auriscalpium (Hydnum), 477
*australiense* (Hydnangium), 463
**australiensis** (Arcangeliella), 463
*australiensis* (*Octaviana*), 463

## B

Bacteria, effect of ultra-violet radiation on, 18, 505
badio-ferruginea (Hymenochaete), 483
Beardsley, Martha L. The cytology of Funaria flavicans Michx. with special reference to fertilization, 509
bellus (Helicomyces), 10, 16
benzoinus (Polyporus), 471
Berkley, Earl E. Studies of the effects of different lengths of day, with variations in temperature, on vegetative growth and reproduction in cotton, 573
bicolor (Odontia), 479
biformis (Polystictus), 471
*Bjerkandera adusta*, 339, 340
Blepharodon **minimus**, 557; suberectus, 557
*Boletus abietinus*, 328; *adustus* 339, var. *crispus*, 340; *albidus*, 337; *applanatus*, 374; *arcularius*, 346, 394; *benzoinus*, 338; *caesius*, 337; *calceolus*, 358; *carpineus*, 339; *cinnabarinus*, 342; *cinnamomeus*, 354; *coccineus*, 342; *conchatus*, 384; *confluens*, 352, 353; *coriaceus*, 353; *crispus*, 340; *elegans*, 359; *ferruginosus*, 401; *fuliginosus*, 338; *fulvus*, 377, 383; *fuscoporus*, 339; *hirsutus*, 324; *igniarius*, 382; *incarnatus*, 328; *incertus*, 397; *isabellinus*, 339; *lipsiensis*, 374; *marginatus*, 377; *medulla-panis*, 398; *nigromarginatus*, 324; *nummularis*, 359; *odoratus*, 370; *ovinus*, 350; *pelleporus*, 339; *perennis*, 353; *Pini*, 379; *pinicola*, 377; *pomaceus*, 383; *resinosus*, 338; *Rheades*, 336; *roseus*, 387; *salicinus*, 384; *semiovatus*, 377; *sistotremoides*, 347; *spumeus*, 345; *squamosus*, 356; *suberosus flabelliformis*, 339; *subtomentosus*, 353; *ungulatus*, 377; *varius*, 358; *versicolor*, 323
*brisbanense* (*Hydnangium*), 463
brumalis (Polyporus), 471
Bucholtz, Alexander F. The effect of monochromatic ultra-violet light of

measured intensities on behavior of plant cells, 489
Burt, Edward Angus. Hymenomycetous fungi of Siberia and eastern Asia—mostly of wood-destroying species, 469
byssoidea (Coniophora), 485

## C

caesio-alba (Poria), 476
calcea (Sebacina), 487
Calcium content of higher plants, effect of ultra-violet radiation on, 565
Callirhoë, 125; *epicata*, 159, 165
Calyptospora columnaris, 293
*Cantharellus alveolaris*, 394
*Capnorea pumila*, 286
Card, H. H. A revision of the genus Frasera, 245
Carestianum (Septobasidium), 487
*Celtidis f. fulvescens* (*Gyroceras*), 4
*Celtidis* (*Gyroceras*), 2, 3
**Celtidis** (Helicoceras), 3, *8*
*Celtidis* (*Monilia*), 3
Chailletii (Stereum), 480
Chromosome complements of Allium stellatum and Nothoscordum bivalve, 465
**cinnamomeus** (Alpova), 461, *464*
citrinus (Leucogaster), 462
*Cladomeris Schweinitzii*, 347
Clarksville soils in Missouri, 48
Clavaria formosa, 486
Clavariaceae, 486
*colliculosum* (*Corticium*), 484
Colorado: The Polyporaceae of, 287; zones of vegetation in, 289
*Coltricia cinnamomea*, 354; *connata*, 353; *perennis*, 353; *tomentosa*, 349
commune (Schizophyllum), 470
confluens, (Corticium), 483
conformis (Ditiola), 486
confragosa (Daedalea), 475
Coniophora arida, 485; byssoidea, 485; olivacea, 485; **sibirica**, 485
Coremiella saxonicum, 7
*Coriolellus Sepium*, 366; *serialis*, 365
*Coriolus abietinus*, 328; *hexagoniformis*, 366; *hirsutulus*, 323; *Lloydii*, 326; *nigromarginatus*, 324; *pargamenus*, 330; *planellus*, 344; *prolificans*, 330; *subchartaceus*, 331; *versicolor*, 323; *zonatus*, 325
cornucopioides (Craterellus), 479
corticalis (Peniophora), 484
Corticium *colliculosum*, 484; confluens, 483; frustulosum, 483; galactinum, 483; hydnans, 484; illaqueatum, 484; investiens, 484; lactescens, 484; laeve, 484; ochraceum, 484; polygonium, 484; radiosum, 484; roseum, 484; sulphureum, 484
Cotton, Studies of the effects of different lengths of day, with variations in temperature, on vegetative growth and reproduction in, 573
Craterellus cornucopioides, 479
*Cryptoporus volvatus*, 321
Cucumbers, effect of ultra-violet radiation: on growth, 27, 30, 568; on calcium content, 570; on phosphorus content, 569
Curtisii (Helicoma), 10
Cyanide, effect of, on cultures of Fusarium lini, 75
Cyclomyces, 314
Cytidia salicina, 486
Cytokinesis in the Mosses, 533
Cytology of Funaria flavicans Michx. with special reference to fertilization, 509, *536*, *538*, *540*

## D

Dacryomycetaceae, 486
Dacryomyces abietinus, 292
Daedalea, 314, 376; aurea, 476; confragosa, 475; *epigaea*, 347; *heteromorpha*, 366; *mollis*, 367; *Pini*, 379; quercina, 476; *saepiaria*, 391; *trabea*, 392; unicolor, 476, var. *violacea*, 328
Dasyscypha Agassizii, 292; arida, 292
Day, effects of different lengths of, with variations in temperature, on vegetative growth and reproduction in cotton, 573
delectans (Polyporus), 472
De Winton, Dorothea, Edgar Anderson and. The genetic analysis of an unusual relationship between self-sterility and self-fertility in Nicotiana, 97
dichrous (Polyporus), 472
*diffissa* (*Peniophora*), 485
**diffissus** (Aleurodiscus), 485
Dipladenia **Achrestogyne**, 543; congesta, 544; fragrans, 545; glabra, 545; Martiana, 545; **oblongifolia**, 544; surinamensis, 545; **upatae**, 545; urophylla, 545
disciformis (Aleurodiscus), 485
Distribution of some Rocky Mountain plants, 283
Ditiola conformis, 486
Dodecatheon tetrandrum, 284
Dodge, C. W. **Alpova**, a new genus of Rhizopogonaceae, with further notes on Leucogaster and Arcangeliella, 457
dryadeus (Polyporus), 472

## E

Echites *agglutinata*, 552; *circinalis*, 552; *coalita*, 552; *cognata*, 546; *Dusenii*,

552; *portobellensis*, 553; *Sanctae-Martae*, 555; *spectabilis*, 552
Eichleriella spinulosa, 487
*Elfvingia applanata*, 374; *lipsiensis*, 374; *megaloma*, 374
ellipticum (Helicoon), 11
*Elytropus spectabilis*, 552
Erinaceus (Hydnum), 477
Exidia glandulosa, 292, 487
Exolobus **marmoreus**, 561; patens, 562

F

Farlowii (Podaxis), 459, *464*
fasciatum (Stereum), 480
Favolus, 394; alveolaris, 394, *454; canadensis*, 394; *europaeus*, 394; *ohiensis*, 394; *striatulus*, 394
Femsjonia luteo-alba, 487
Fertilization: in Funaria flavicans Michx., 509, *536*, *538*, *540*; in the Mosses, 509; in the Liverworts, 511
fibrillosus (Polyporus), 472
Flax, glucoside content of, 66
floccosus (Leucogaster), 459
Fomes, 376; *Abietis*, 379; *alboluteus*, 333; annosus, 389, *450; applanatus*, 374; conchatus, 384; connatus, 396; Demidoffii, 385, 388, *446; Earlei*, 385; *Ellisianus*, 388; fraxinophilus, 389, f. Ellisianus, 388, *452; fuliginosus*, 338; fulvus, 383, *446*, 474; igniarius, 382, *448*, 474; *Juniperinus*, 385; *leucophaeus*, 374; *marginatus*, 377; *megaloma*, 374; *nigricans*, 382; nigrolimitatus, 380, *454;* ohiensis, 376; Palliseri, 474; Pini, 379, *442;* pinicola, 377, *444*, *446*, 474; *pomaceus*, 383; *ponderosus*, 378; putearius, 381; roseus, 372, 387, *436*, 474; scutellatus, 376; *tenuis*, 363; *ungulatus*, 377
Fomitopsis *annosa*, 389; *pinicola*, 377; *rosea*, 387
formosa (Clavaria), 486
Fossombronia, fertilization in, 512, 530
Frasera, A revision of the genus, 245
Frasera, 256; *albicaulis*, 269, f. *alba*, 269; albomarginata, 274, *280*, var. **induta**, 275; *ampla*, 259; *angustifolia*, 259; *carolinensis*, 262; caroliniensis, 262, *280;* coerulea, 271; *Cusickii*, 270; fastigiata, 264, *280; induta*, 275; *macrophylla*, 259; montana, 266, *280;* neglecta, 265, *280;* nitida, 267, *280*, var. **albicaulis**, 269, *280*, var. **Cusickii**, 270, *280*, *282; nitida* var. *albida*, 267; *officinalis*, 262; paniculata, 275, *280;* Parryi, 273, *280;* puberulenta, 272, *280*, *282;* speciosa, 258, *280; speciosa* var. *angustifolia*, 259, var. *scabra*, 259, var. *stenosepala*, 259; *stenosepala*, 259; *thyrsiflora*, 264; tubulosa, 271, *280; utahensis*, 275; *venosa*, 259; *verticillata*, 262; *Walteri*, 262
Friesii (Porothelium), 476
frondosus (Polyporus), 472
frustulosum (Corticium), 483
Fuller, Harry J. Stimulatory effects of radiation from a quartz mercury vapor arc upon higher plants, 17; F. Lyle Wynd and. Some effects of ultra-violet radiation upon the calcium and phosphorus content of higher plants, 565
*Fulvifomes Juniperinus*, 385
fulvus (Fomes), 474
*Funalia stuppea*, 368
Funaria flavicans Michx., cytology of, 509; hygrometrica, 512
Fusarium lini, 66
**fuscopes** (Helicomyces), 15
*Fuscoporia ferruginosa*, 401
fusco-violaceus (Irpex), 478
fuscum (Stereum), 481

G

galactinum (Corticium), 483
Ganoderma, 373; applanatum, 374, *442; leucophaeum*, 374; *lipsiensis*, 374
Gasteromycetes, 457
gausapatum (Stereum), 481
Genetic analysis of an unusual relationship between self-sterility and self-fertility on Nicotiana, 97
gibbosa (Trametes), 475
gigantea (Peniophora), 485
gilvus (Polyporus), 472
*glabrum* (Hydnangium), 463
glandulosa (Exidia), 487
*Gloeophyllum abietinellum*, 391; *hirsutum*, 370, 391; *pallidofulvum*, 393; *trabeum*, 393
Glucoside content of flax, 66
Goodman, George J. Notes on the distribution of some Rocky Mountain plants, 283
griseum (Helicosporium), 9
Guepinia monticola, 292
*Gyroceras*, 1; *ammonis*, 1, 7; *Celtidis*, 2, 3, f. *fulvescens*, 4; *divergens*, 4; *Nymphaearum*, 5; *Plantaginis*, 5; *sazonicum*, 7
Gyrocerus, 2

H

*Haemadictyon circinalis*, 552
Hagerstown soils in Missouri, 44
Helicoceras, The genus, 1
**Helicoceras**, 1; **Celtidis**, 3, *8;* **Nymphaearum**, 5, *8;* **Oryzae**, 2, 6, *8;* **Plantaginis**, 5, *8*
Helicoma ambiens, 10; **anastomosans**, 13, 14, *16;* Curtisii, 10; Morgani, 13, 14; olivaceum, 11; polysporum, 11; repens,

11; **tenuifilum**, 14, *16;* violaceum, 14; **Westoni**, 12, *16*
Helicomyces bellus, 10, 16; **fuscopes**, 15; roseus, 10, 16
Helicoon ellipticum, 11
Helicosporeae, Brief notes on the, with descriptions of four new species, 9
Helicosporium griseum, 9; lumbricoides, 10; lumbricopsis, 10; *Nymphaearum*, 5; phragmites, 9
*Hesperalcea*, 137, 214; *malachroides*, 128, 215
Hesperochiron pumilus, 284
*Heterobasidion annosum*, 389
heteromorpha (Lenzites), 470
*Hexagona*, 394; *alveolaris*, 394; *micropora*, 394; *Mori*, 394; *striatula*, 394
hirsutum (Stereum), 481
hirsutus (Polystictus), 471
hispida (Trametes), 474
hispidus (Polyporus), 472
Hollii (Hydnum), 477
Host plants for species of Polyporaceae, 294
Hydnaceae, 477
*Hydnangium australiense*, 463; *brisbanense*, 463; *glabrum*, 463; *nudum*, 462; *Ravenelii*, 463; *Stephensii*, 463, v. *Ravenelii*, 463; *virescens*, 462
hydnans (Corticium), 484
Hydnum auriscalpium, 477; Erinaceus, 477; Hollii, 477; **Murashkinskyi**, 477; ochraceum, 478; **reflexum**, 478; velutinum, 478
Hydrocyanic acid in flax, 66
Hymenochaete badio-ferruginea, 483; Mougeotii, 483; tabacina, 483
Hymenomycetous fungi of Siberia and eastern Asia—mostly of wood-destroying species, 469
Hypochnus spongiosus, 480; umbrinus, 480
Hysterangium, 457

## I

igniarius (Fomes), 474
illaqueatum (Corticium), 484
inodora (Trametes), 474
*Inonotus dryophilus*, 336; *fibrillosus*, 334
investiens (Corticium), 484
Irpex, 334; fusco-violaceus, 478; *grossus*, 368; lacteus, 478; pachyodon, 479
*Ischnoderma fuliginosum*, 338; *resinosum*, 338

## J

Juglandis (Microstroma), 486

## K

Kneiffia setigera, 479

## L

lacteus (Irpex), 478
lacteus (Polyporus), 473
lactescens (Corticium), 484
laevigata (Poria), 476
laricina (Lenzites), 470
Laubertia Boissieri, 556; **Pringlei**, 555; **Sanctae-Martae**, 555
laeve (Corticium), 484
lentus (Polyporus), 473
Lenzites, 390; abietina, 392; *abietinella*, 391, *452; abietis*, 328; heteromorpha, 470; *heteromorpha*, 366, 470; laricina, 470; *rhabarbarina*, 391; saepiaria, 391, *452; saepiaria porosa*, 370; septentrionalis, 470; serpens, 315; tenuis, 470; trabea, 392, *454; vialis*, 392
Lesquerella Kingii, 284; prostrata, 284
*Leucocraspedum*, 257; *albicaule*, 269; *albomarginatum*, 274; *coeruleum*, 271; *montanum*, 266
Leucogaster, 457; citrinus, 462; floccosus, 459; *luteomaculatus*, 462; nudus, 462
Light: effect of, on glucoside content of flax, 73; effect of, on growth and reproduction in plants, 573; ultra-violet, effect of measured intensities on behavior of plant cells, 489
Linamarine, 90
Linder, D. H. The genus Helicoceras, 1; Brief notes on the Helicosporeae, with descriptions of four new species, 9
Linkage between self-sterility allelomorphs and other genes, 108
Liverworts, fertilization in, 511
lumbricoides (Helicosporium), 10
lumbricopsis (Helicosporium), 10
luteo-alba (Femsjonia), 487
*luteomaculatus* (*Leucogaster*), 462

## M

Macropharynx *fistulosa*, 552; **spectabilis**, 552
Macroscepis barbata, 560; **equatorialis**, 559
Malouetia **Killipii**, 551; pubescens, 552
Malva, 125; *malachroides*, 215, 217
Malvaceae, 124
Malvastrum 128; *confertum*, 213; spicatum, 128
Mandevilla, 556
Marsdenia **lauretiana**, 562; mollissima, 563
Melanogaster, 457
melanopus (Polyporus), 473
Mertensia incongruens, 286
*Merulius*, 322; *alveolaris*, 394; serpens, 477; tremellosus, 477
*Mesadenia*, 246, 257
Microstroma Juglandis, 486

Missouri soils, acidity of, 41
Mnium: effect of ultra-violet radiation on cell contents of leaves, 500; fertilization in, 510
Moisture: influence of, upon spore germination and growth, 300; soil, effect of, on growth of cotton plants, 578
Monilia Celtidis (Celtis), 2
*Monilia Celtidis*, 2, 3
Monochromatic ultra-violet light of measured intensities, Effect of, on behavior of plant cells, 489
Monochromator in ultra-violet radiation work, 496
Morgani (Helicoma), 13, 14
Mosses, fertilization in, 509
mucida (Poria), 476
Murashkinsky, K. E. Fungi received from, 469
**Murashkinskyi** (Hydnum), 477
Mougeotii (Hymenochaete), 483
mutata (Peniophora), 485
*Myriadoporus adustus*, 339

## N

Nicotiana, self-sterility and self-fertility in, 97
Nicotiana alata, 97, × Langsdorfii, 97; Sanderae, 97
Nothoscordum bivalve, Chromosome complements of, 465
*nudum* (*Hydnangium*), 462
nudus (Leucogaster), 462
*Nuttallia malvaeflora*, 188
*Nymphaearum* (*Gyroceras*), 5
**Nymphaearum** (Helicoceras), 5, *8*
*Nymphaearum* (*Helicosporium*), 5

## O

ochraceum (Corticium), 484
ochraceum (Hydnum), 478
ochroleucum (Stereum), 469, 481
*Ochroporus odoratus*, 370
*Octaviania australiensis*, 463; *Ravenelii*, 463; *rosea*, 462; *Stephensii*, 463, v. *Ravenelii*, 463
*Octavianina Stephensii*, 463
Odontadenia **affinis**, 549; **augusta**, 548; **cognata**, 546; coriacea, 551; Cururu, 549; geminata, 551; **glauca**, 550; hypoglauca, 551; **Killipii**, 546; nitida, 551; **Perrottetii**, 546; **polyneura**, 546, 551; **Sandwithiana**, 547; speciosa, 548; **stemmadeniaefolia**, 548
Odontia bicolor, 479
olivacea (Coniophora), 485
olivaceum (Helicoma), 11
**Oryzae** (Helicoceras), 6, *8*
osseus (Polyporus), 473

## P

pachyodon (Irpex), 479
Palliseri (Fomes), 474
palmata (Thelephora), 479
Paramecia, effect of ultra-violet radiation on, 505
Paxina nigrella, 293
Pellia, fertilization in, 512, 530
*Pelloporus perennis*, 353
Peniophora corticalis, 484; *diffissa*, 485; gigantea, 485; mutata, 485; *quercina*, 484
pergamenus (Polystictus), 471
*Phaeoporus applanatus*, 374
*Phaeolis sistotremoides*, 347
Phaeostemma grandifolia, 561; **tigrina**, 560
*Phellinus igniarius*, 382; *isabellinus*, 363; *nigrolimitatus*, 380; *salicinus*, 384
Phellorinia, 459
Phlebia radiata, 479; strigoso-zonata, 479
Phosphorus content of higher plants, effect of ultra-violet radiation on, 565
phragmites (Helicosporium), 9
Physiology of plant disease, Studies in the, 57
Pini (Stereum), 482
pinicola (Fomes), 474
*Placodes applanatus*, 374; *fulvus*, 383; *pomaceus*, 383
*Plantaginis* (*Gyroceras*), 5
**Plantaginis** (Helicoceras), 5, *8*
*plantaginis* (Torula), 5
Pleurococcus, effect of ultra-violet radiation on cell contents of, 499
Pleurotus applicatus, 470
Podaxis, 459; anomalus, 459; Farlowii, 459, *464*
polygonium (Corticium), 484
Polyporaceae, 470; of Colorado, 287; factors determining the distribution of, 296
Polyporus, 317; abietinus, 328, *414;* adustus, 339, *420*, 471; alboluteus, 333, *416;* amorphus, 471; anceps, 343, *424; annosus*, 389; *applanatus*, 374; *arculariformis*, 346; arcularius, 346, *426; aurantiacus*, 334; benzoinus, 471; *benzoinus*, 338; borealis, 335, *418; Boucheanus peponinus*, 394; brumalis, 471; caesius, 337, *434; calceolus*, 358; *callosus*, 365; *caudicinus*, 356; *cervinus*, 367; cinnabarinus, 342, *426;* cinnamomeus, 354, *426;* circinatus, 349, *430; circumstans*, 388; *conchatus*, 384; confluens, 351, *434; corruscans*, 336; crispus, 340, *424;* cryptopus, 355, *428;* delectans, 472; *Demidoffii*, 385; dichrous, 472; dryadeus, 472; *dryophilus*, 336; *dualis*, 349; elegans, 359, *432;*

Ellisianus, 343; *ferruginosus*, 401; fibrillosus, 316, 334, *412*, 472; *Flabellum*, 329; fragilis, 341, *426;* frondosus, 472; *fuliginosus*, 338; *fulvus*, 338; *Gillotii*, 389; gilvus, 472; *hirsutulus*, 323; hirsutus, 324, *412;* hirtus, 357, *432; hispidellus*, 357; *hispidoides*, 347; hispidus, 472; *hyperboreus*, 382; *igniarius*, 382; *ilicincola*, 330; *inflatus*, 321; *irregularis*, 389; *Juniperinus*, 385; *laceratus*, 329; lacteus, 473; *laetificus*, 400; lentus, 473; leucomelas, 288; *leucophaeus*, 374; leucospongia, 326, *412; Lindheimeri*, 368; *Lloydii*, 326; *loricatus*, 384; *Macouni*, 401; *marginatus*, 377; *medulla-panis*, 398; *megaloma*, 374; melanopus, 473; *Menandianus*, 329; mollis, 342; *nigricans*, 382; *nigrolimitatus*, 380; *Novae-Angliae*, 382; *oblectans*, 354; *obvolutus*, 321; *odoratus*, 370; osseus, 360, 473; ovinus, 350, *426; pallescens*, 339; pargamenus, 329, *410; parvulus*, 328, 354; *Peakensis*, 349; perennis, 353, *430; piceinus*, 379; picipes, 359; *Pini*, 379; *pinicola*, 377; planellus, 344, *422; planus*, 344; *pomaceus*, 383; *prolificans*, 329; *pseudopargamenus*, 330; pubescens, 326; *purpurea*, 401; resinosus, 338, 389, *422*, 473; Rheades, 336, *420; roseus*, 387; *rufo-pallidus*, 387; *salicinus*, 384; *salmonicolor*, 400; *Sartwellii*, 330; *scalaris*, 365; Schweinitzii, 347, *428; serialis*, 365; *serpentarius*, 389; *Shiraianus*, 334; *spectabilis*, 347; *spissus*, 400; splendens, 354; spumeus, 345, *434*, 473; squamosus, 356, 473; *steroides*, 367; *subacida*, 399; subchartaceus, 331, *414; subcinereus*, 339; *subflavus*, 330; *subpileatus*, 389; *subsericeus*, 354; subsquamosus, 356; *tabulaeformis*, 347; *tenuis*, 363; *tomentosus*, 349; trichrous, 473; *ursinus*, 332, *416; vaporaria*, 397; *variiformis*, 365; varius, 358, *432;* versicolor, 323, *410;* volvatus, 321, *410; volvatus Helix*, 321; *xalapensis*, 330; *xantholoma*, 398; *Zelleri*, 360; zonatus, 325, *410*

polysporum (Helicoma), 11

*Polystictus*, 313; *abietinus*, 328, 470; *azureus*, 323; *biformis*, 471; *Celottianus*, 368; *cinnabarinus*, 342; *cinnamomeus*, 354; *Fergussoni*, 368; *hirsutus*, 324, 471; *hirtellus*, 324; *pargamenus*, (*pergamenus*), 330; *perennis*, 353; pergamenus, 471; *piceinus*, 379; *pusio*, 328; radiatus, 471; *Schweinitzii*, 347; *scuirinus*, 368; *subchartaceus*, 331; *versicolor*, 323; vulpinus, 471; *zonatus*, 326

Poria, 313, 395; caesio-alba, 476; *callosa*, 365, 396; contigua, 364; ferruginosa, 364, 401; *incerta*, 397; *laetifica*, 400; laevigata, 476; *Macouni*, 401; medullapanis, 398, *456;* monticola, 396, *454;* mucida, 398, 476; *obducens*, 396; purpurea, 401; *salmonicolor*, 400; spissa, 400, *456;* subacida, 399, *456;* superficialis, 364; taxicola, 476; Vaillantii, 398; vaporaria, 397, *456;* violacea, 401; viticola, 364; xantha, 476; *xantholoma*, 398

*Porodaedalea Pini*, 379

Porothelium Friesii, 476

Prestonia acutifolia, 554; **agglutinata**, 552; **coalita**, 552; **Dusenii**, 552; **isthmica**, 555; *Langlassei*, 555; *macrocarpa*, 553; mexicana, 555; Muelleri, 553; **portobellensis**, 553; Riedelii, 553; **velutina**, 554

protracta (Trametes), 475

pubescens (Polyporus), 473

*Pycnoporellus fibrillosus*, 334

*Pycnoporus cinnabarinus*, 342

*Pyropolyporus conchatus*, 384; *Earlei*, 385; *fulvus*, 383; *Juniperinus*, 385

## Q

Quartz mercury vapor arc, Stimulatory effects of radiation from, 17

quercina (Daedalea), 476

*quercina* (*Peniophora*), 484

Quinhydrone electrode method for testing soils, 41

## R

radiata (Phlebia), 479

**radiata** (Trametes), 475

Radiation: Stimulatory effects of, from a quartz mercury vapor arc, 17; ultra-violet, effect of, upon plants, 489, upon the calcium and phosphorus content of higher plants, 565

radiatus (Polystictus), 471

radiosum (Corticium), 484

Rauwolfia bahiensis, 543; **lauretiana**, 541; paucifolia, 542; **sanctorum**, 542

**Ravenelii** (Arcangeliella), 463

*Ravenelii* (*Hydnangium*), 463

*Ravenelii* (*Octaviania*), 463

**reflexum** (Hydnum), 478

repens (Helicoma), 11

resinosus (Polyporus), 473

Reynolds, Ernest S. Studies on the physiology of plant disease, 57

*Rhabdadenia polyneura*, 546

*Rhaptocarpus coalitus*, 552

Rhizopogon, 457

Rhoeo discolor, effects of ultra-violet radiation on cell contents of, 499

rhytidocyclum (Stereum), 482

Riccardia, fertilization in, 512, 530
Rocky Mountain plants, Notes on the distribution of some, 283
*rosea* (*Octaviania*), 462
**rosea** (Arcangeliella), 462
roseum (Corticium), 484
roseus (Fomes), 474
roseus (Helicomyces), 10, 16
Roush, Eva M. Fling. A monograph of the genus Sidalcea, 117
rugosiusculum (Stereum), 482
rugosum (Stereum), 482
rutilans (Polyporus), 473

## S

salicina (Cytidia), 486
sanguinolentum (Stereum), 482
Sarcopodium, 2
Sarcopodium (Tricholeconium) **ammonis**, 7
saxonicum (Coremiella), 7
*saxonicum* (*Gyroceras*), 7
Schizophyllum commune, 470
*Scutiger cryptopus*, 355; *hispidellus*, 357; *laeticolor*, 352; *ovinus*, 350; *Whiteae*, 352
Sebacina calcea, 487
Self-sterility and self-fertility in Nicotiana, Genetic analysis of an unusual relationship between, 97
septentrionalis (Lenzites), 470
Septobasidium Carestianum, 487
serpens (Merulius), 477
*Sesia hirsuta*, 391; *pallidofulva*, 393
setigera (Kneiffia), 479
Shope, Paul Franklin. The Polyporaceae of Colorado, 287
Siberia, Hymenomycetous fungi of, 469
**sibirica** (Coniophora), 485
Sida, 117, 125, 197; *californica*, 199; *delphinifolia*, 143, 145, 188, 193, 197; *diploscypha*, 139, *heterophylla*, 173; *malvaeflora*, 175, 188; *oregana*, 168
Sidalcea, A monograph of the genus, 117
Sidalcea, 117, 136; sect. **Annuae**, 138; *asplenifolia*, 175, 177; asprella, 201; *atacosa*, 218; atropurpurea, 218; *californica*, 199; calycosa, 145, *230;* campestris, 175; *campestris*, 169, 218; candida, 153, *228*, *234; candida* var. *tincta*, 153; *confinis*, 179; *Covillei*, 187; *crenulata*, 179; *Cusickii*, 174; *delphinifolia*, 188, var. *humilis*, 189; diploscypha, 139, *228*, *232; diploscypha* var. *minor*, 139; *elegans*, 202; subgenus Eusidalcea, 129, 138; *eximia*, 159; *favosa*, 200; glaucescens, 207; Hartwegi, 148, *230; Hartwegi* var. *tenella*, 148; subgenus Hesperalcea, 128, 214; Hendersoni, 157; Hickmani, 212; Hickmani var. Parishii, 213, *228*, *230*, *244; Hickmani*, 213; hirsuta, 143, *230; humilis*, 188; *hydrophila*, 159; *interrupta*, 166; malachroides, 215, *228*, *230*, *244;* malvaeflora, 188, *228*, *240*, var. atropurpurea, 218, var. californica, 199, var. Listeri, 218; *malvaeflora*, 157, 168, 179, var. *asprella*, 202, var. *Oregana*, 157, 168; subgenus **Malvastralcea**, 128, 212; mariana, 218; mexicana, 218; *montana*, 207; multifida, 209, *240; "Murrayana,"* 159; *Nelsoniana*, 159; neo-mexicana, 179, var. **Covillei**, 187, var. **parviflora**, 186; *neo-mexicana* var. *Diehlii*, 179; *nervata*, 169; *nitrophila*, 186; nodosa, 218; oregana, 168, 177, *230*, *236*, var. **Cusickii**, 174; *oregana*, 175; *Parishii*, 213; *parviflora*, 186, var. *Thurberi*, 179; pedata, 210, *242;* sect. **Perennes**, 151; peruviana, 218; *ranunculacea*, 166; reptans, 200; *rhizomata*, 145; **robusta**, 205; *rostrata*, 189; *scabra*, 189; *secundiflora*, 139; spicata, 159, var. **ranunculacea**, 166, *230*, *238; spicata* var. *pedata*, 210, var. *reptans*, 200; *sulcata*, 145; *sylvestris*, 175; *tenella*, 148; triloba, 218; *valida*, 159; virgata, 178; *vitifolia*, 215
Sidalcia, 145
Soils in Missouri, the acidity of, and its relation to plant distribution, 41
spadiceum (Stereum), 483
Sphaeralcea, 213; pedatifida, 218
Sphaerocarpos, fertilization in, 511, 530
spinulosa (Eichleriella), 487
*Spongipellis borealis*, 335; *fragilis*, 341; *occidentalis*, 345; *spumeus*, 345
spongiosus (Hypochnus), 480
*Spongiporus leucospongia*, 326
spumeus (Polyporus), 473
squamosus (Polyporus), 473
Stephanotella Glaziovii, 558; **Killipii**, 558
**Stephensii** (Arcangeliella), 463
*Stephensii* (*Hydnangium*), 463
*Stephensii* (*Octaviania*), 463
*Stephensii* v. *Ravenelii* (*Hydnangium*), 463
*Stephensii* (*Octavianina*), 463
stereoides (Trametes), 475
Stereum bicolor, 481; Chailletii, 480; fasciatum, 480; fuscum, 481; gausapatum, 481; hirsutum, 481; ochroleucum, 469, 481; Pini, 482; rhytidocyclum, 482; rugosiusculum, 482; rugisporum, 292; rugosum, 482; sanguinolentum, 482; spadiceum, 483; sulcatum, 483; versiforme, 483
Steyermark, Julian A. A study of plant distribution in relation to the acidity of various soils in Missouri, 41

Stimulatory effects of radiation from a quartz mercury vapor arc upon higher plants, 17
*Streptotrachelus Pringlei*, 555
*Strilia cinnamomea*, 354
Studies on the physiology of plant disease, 57
suaveolens (Trametes), 475
sulcatum (Stereum), 483
sulphureum (Corticium), 484
*Swertia*, 257; *albicaulis*, 269; *albomarginata*, 274; *Bigelowii*, 279; *carolinensis*, 262; *fastigiata*, 264; *Parryi*, 273; *radiata*, 259
Swertia, 245, 247
*Swertia*, 257; *albomarginata*, 274; *difformis*, 262; *fastigiata*, 264; *Frasera*, 262; *neglecta*, 263; *nitida*, 265; *Parryi*, 273; *tubulosa*, 271

T

tabacina (Hymenochaete), 483
Tabernaemontana hirtula, 541; **Killipii**, 541
taxicola (Poria), 476
Temperature: effect of, on glucoside content of flax, 73; effect of variations in, and lengths of day, on vegetative growth and reproduction in cotton, 573; influence of, upon spore germination and growth, 297
**tenuifilum** (Helicoma), 14, *16*
tenuis (Lenzites), 470
**tenuis** (Thelephora), 479
terrestris (Thelephora), 480
*Tessaranthium*, 257; *angustifolium*, 259; *macrophyllum*, 259; *scabrum*, 259; *speciosum*, 259; *stenosepalum*, 259
*Tesseranthium*, 246, 257; *radiatum*, 259
Thelephora palmata, 479; **tenuis**, 479; terrestris, 480
Thelephoraceae, 479
Tilsit soils in Missouri, 42
Tomatoes, effect of ultra-violet radiation: on growth, 28, 36, 568; on calcium content, 570; on phosphorus content, 569
*Torula Plantaginis*, 5
Tradescantia reflexa, effect of ultra-violet radiation on cell contents of stamen hairs, 499
Trametes, 362; Abietis, 474; *benzoina*, 338; *carnea*, 372; *cinnabarina*, 342; gallica, 369; gibbosa, 475; heteromorpha, 366, *440;* hispida, 368, *440*, 474; inodora, 474; isabellina, 363, *438; laceratus*, 366; *mollis*, 367; odorata, 370, *440; Peckii*, 368; *Pini*, 379; *Pini Abietis*, 379; *protracta*, 370, 475; **radiata**, 475; *radiciperda*, 389; serialis, 365, *438; setosus*, 363; stereoides, 367, *440*, var. Kmetii, 475; stuppeus, 370; suaveolens, 475; subrosea, 372, *442; tenuis*, 363; *trabea*, 475; variiformis, 365, *438*
tremellosus (Merulius), 477
Tremellaceae, 487
Tricholeconium, 2
trichrous (Polyporus), 473
*Tyromyces anceps*, 343; *caesius*, 337; *Ellisianus*, 343

U

Ultra-violet light, Effect of monochromatic, of measured intensities, on behavior of plant cells, 489
Ultra-violet radiation: effects of, upon plants, 17, 489; Some effects of, upon calcium and phosphorus content of higher plants, 565
umbrinus (Hypochnus), 480
*Ungulina annosa*, 389; *fuliginosa*, 338; *marginata*, 377; *rosea*, 387; *volvata*, 321, var. *pleurostoma*, 321
unicolor (Daedalea), 476
Union soils in Missouri, 47

V

velutinum (Hydnum), 478
versiforme (Stereum), 483
*Villarsia pumila*, 286
violaceum (Helicoma), 14
*virescens* (*Hydnangium*), 462
Vita-glass filter in ultra-violet radiation experiments, 30
vulpinus (Polystictus), 471

W

**Westoni** (Helicoma), 12, *16*
Wherry's soil acidity results, compared with Steyermark's, 51
Woodson, Robert E. Jr. New or otherwise noteworthy Apocynaceae of tropical America, 541; New South American Asclepiadaceae, 557
Wynd, F. Lyle, and Harry J. Fuller. Some effects of ultra-violet radiation upon the calcium and phosphorus content of higher plants, 565

X

xantha (Poria), 476

Z

Zea Mays, effect of ultra-violet radiation on, 566